

Artin HNN-extensions virtually embed in Artin groups

T. Hsu* I. J. Leary*[†]

February 12, 2008

Abstract

An Artin HNN-extension is an HNN-extension of an Artin group in which the stable letter conjugates a pair of suitably chosen subsets of the standard generating set. We show that some finite index subgroup of an Artin HNN-extension embeds in an Artin group. We also obtain an analogous result for Coxeter groups.

1 Results

Recall that an *Artin system* is a finite set S , together with a function m from the 2-element subsets of S to the set $\{2, 3, 4, \dots\} \cup \{\infty\}$. It may be helpful to consider m as an edge-labelling on the complete graph with vertex set S . The *Artin group* $A = A(S, m)$ corresponding to an Artin system (S, m) is defined by a presentation. The generators for A are the elements of the set S . There is one relation in the presentation for each unordered pair $s, t \in S$ such that $m(t, s) = m(s, t) < \infty$, of the form

$$sts \cdots = tst \cdots, \quad (1)$$

where there are $m(s, t)$ symbols on each side of (1). Call these relations the *Artin relations* associated to (S, m) . If S' is a subset of S , and we write m for the restriction to S' of m , inclusion induces a group homomorphism from $A(S', m)$ to $A(S, m)$, and this homomorphism is known to be injective [8, 9]. The image of such a homomorphism is called an *Artin subgroup* of $A(S, m)$.

An Artin system also gives rise to a *Coxeter group* $W = W(S, m)$. The generators of the Coxeter group are the elements of S , subject to the Artin relations for (S, m) together with the relations $s^2 = 1$ for all $s \in S$. Call these relations the *Coxeter relations* for W . Note that in the presence of the relations $s^2 = 1 = t^2$, the Artin relation between s and t is equivalent to the relation

*This work was done at MSRI, Berkeley, where research is supported in part by NSF grant DMS-0441170

[†]partially supported by NSF grant DMS-0505471.

$(st)^{m(s,t)} = 1$. For S' a subset of S , the natural map $W(S', m) \rightarrow W(S, m)$ is injective, and the image of such a homomorphism is called a *Coxeter subgroup* of $W(S, m)$ [1, 2].

The *set of labels*, $L(S, m)$, for an Artin system (S, m) is the image of m , viewed as a subset of $\{2, 3, 4, \dots\} \cup \{\infty\}$. An Artin system (S, m) is called *right-angled* if $L(S, m) \subseteq \{2, \infty\}$, and $A(S, m)$ (resp. $W(S, m)$) is then called a *right-angled Artin group* (resp. *right-angled Coxeter group*).

Now suppose $\phi : S' \rightarrow S''$ is a bijection between two subsets of S that preserves labels in the sense that

$$m(s, s') = m(\phi(s), \phi(s')) \quad \text{for all } s, s' \in S'. \quad (2)$$

The *Artin HNN-extension* $G(S, m, \phi)$ (resp. *Coxeter HNN-extension* $C(S, m, \phi)$) associated to (S, m, ϕ) is the free product of $A(S, m)$ (resp. $W(S, m)$) and an infinite cyclic group $\langle t \rangle$, modulo the relations $t^{-1}st = \phi(s)$ for all $s \in S'$. Thus an Artin HNN-extension (resp. a Coxeter HNN-extension) is a special type of HNN-extension in which the base group is an Artin (resp. a Coxeter) group and the stable letter conjugates an Artin (resp. a Coxeter) subgroup to another Artin (resp. Coxeter) subgroup via the map induced by a bijection of standard generating sets. The *set of labels* for $G(S, m, \phi)$ (resp. $C(S, m, \phi)$) is the set of labels for the group $A(S, m)$ (resp. $W(S, m)$).

We require one more definition before stating our main result. For any property $\mathcal{P}$ of groups, a group G *virtually has property $\mathcal{P}$* if there exists a finite-index subgroup $H \leq G$ such that H has property $\mathcal{P}$.

Theorem 1 *Any Artin (resp. Coxeter) HNN-extension virtually embeds in an Artin (resp. Coxeter) group. If L is the set of labels for the HNN-extension, then the Artin (resp. Coxeter) group in which it virtually embeds can be chosen to have its set of labels equal to $L \cup \{2, \infty\}$. In particular right-angled HNN-extensions virtually embed in right-angled groups.*

Theorem 1 yields the following application, which was our original motivation for considering Artin HNN-extensions.

Corollary 2 *For any finite group Q not of prime power order, there exists n and a subgroup G of $SL(n, \mathbb{Z})$ such that G virtually has a finite classifying space and G has infinitely many conjugacy classes of subgroups isomorphic to Q .*

Proof. Note that by ‘ G virtually has a finite classifying space’ we mean that there is a finite index subgroup $H \leq G$ which has a finite $K(H, 1)$. In [6], a group G is constructed which virtually has a finite classifying space and which contains infinitely many conjugacy classes of subgroup isomorphic to Q . By [6, Thm. 28], we know that G virtually embeds in a right-angled Artin HNN-extension. But then, by Theorem 1, G virtually embeds in a right-angled Artin group, and by [3, 5], any right-angled Artin group embeds in $SL(m, \mathbb{Z})$ for some m , so the group G embeds in $SL(n, \mathbb{Z})$ for some (possibly larger) n . ■

Remark 3 *Corollary 2 was proved in [7] in the special case when Q is not of the form $(p\text{-group})\text{-by-cyclic-by-}(q\text{-group})$ for primes p and q .*

2 Proofs

To review the standard definitions from [10, I.5], recall that a *graph of groups* $(G_{(-)}, \Gamma)$ consists of a directed CW-graph Γ , together with a group G_v for each vertex v of Γ (the *vertex groups*), a group G_e for each edge e of Γ (the *edge groups*), and for each edge e , two injective group homomorphisms $\iota_e : G_e \rightarrow G_{\iota(e)}$ and $\tau_e : G_e \rightarrow G_{\tau(e)}$ (the *attaching homomorphisms*), where $\iota(e)$ denotes the initial vertex of e and $\tau(e)$ denotes the terminal vertex of e . We define $F(G_{(-)}, \Gamma)$ to be the free product of the G_v and infinite cyclic groups $\langle t_e \rangle$, where e runs over all edges of Γ , modulo the relations

$$t_e^{-1} \iota_e(g) t_e = \tau_e(g) \quad \text{for all } e, g \in G_e; \quad (3)$$

and for a basepoint P_0 in Γ , we define $\pi_1(G_{(-)}, \Gamma, P_0)$ to be the subgroup of all words in $F(G_{(-)}, \Gamma)$ that, after ignoring all syllables from G_v 's, define paths in Γ that begin and end at P_0 . Similarly, for a spanning tree T of Γ , we define $\pi_1(G_{(-)}, \Gamma, T)$ to be the free product of the G_v and infinite cyclic groups $\langle t_e \rangle$, where e runs over all edges of Γ not in T , modulo the relations

$$\iota_e(g) = \tau_e(g) \quad \text{for all } e \in T, g \in G_e, \text{ and} \quad (4)$$

$$t_e^{-1} \iota_e(g) t_e = \tau_e(g) \quad \text{for all } e \notin T, g \in G_e. \quad (5)$$

Up to isomorphism, $\pi_1(G_{(-)}, \Gamma, P_0)$ is independent of P_0 and $\pi_1(G_{(-)}, \Gamma, T)$ is independent of T , and the two groups are isomorphic, so we call their isomorphism class the *fundamental group of* $(G_{(-)}, \Gamma)$, denoted by $\pi_1(G_{(-)}, \Gamma)$. We also call the elements t_e *stable letters*.

Let k be a positive integer, let V , E , and V_* be groups, and let $f : E \rightarrow V$, $g : E \rightarrow V$, and $h_i : V \rightarrow V_*$ ($i \in \mathbb{Z}/k$) be injective group homomorphisms.

1. Let Δ_1 be a self-loop. Taking vertex group V , edge group E , and attaching maps f and g , we get a graph of groups $(\{V, E\}, \Delta_1)$ whose fundamental group is an HNN extension with base group V and associated subgroup E . Let t be the associated stable letter.
2. Let Δ_k be a (directed) k -cycle, with vertex set $\mathbb{Z}/k$ and edges $(i, i+1)$ ($i \in \mathbb{Z}/k$). Taking the vertex (resp. edge) groups V_i (resp. E_i) to be copies of V (resp. E), and the attaching homomorphisms $f_i : E_i \rightarrow V_i$ and $g_i : E_i \rightarrow V_{i+1}$ to be copies of f and g , with all indices running over all $i \in \mathbb{Z}/k$, we get a graph of groups $(\{V_i, E_i\}, \Delta_k)$. Let $\{t_i\}$ be the set of stable letters in the group $F(\{V_i, E_i\}, \Delta_k)$.
3. Let Γ_k be a k -leaved rose, with edges labelled by $\mathbb{Z}/k$, and let V_i , E_i , f_i , and g_i be defined as in 2. Taking vertex group V_* , edge groups the E_i , and attaching homomorphisms $h_i \circ f_i$ and $h_{i+1} \circ g_i$, we get a graph of groups $(\{V_*, E_i\}, \Gamma_k)$.

Lemma 4 *Let K be the kernel of the homomorphism $\rho : \pi_1(\{V, E\}, \Delta_1) \rightarrow \mathbb{Z}/k$ defined by $\rho(V) = 0$ and $\rho(t) = 1$. Then $K = \pi_1(\{V_i, E_i\}, \Delta_k)$, and K embeds in $\pi_1(\{V_*, E_i\}, \Gamma_k)$.*

Proof. Let $\theta : \pi_1(\{V_i, E_i\}, \Delta_k, 0) \rightarrow \pi_1(\{V, E\}, \Delta_1)$ be the homomorphism defined by $\theta(v) = v \in V$ for all $v \in V_i$ and $\theta(t_i) = t$. Since θ sends normal forms to normal forms, θ is injective; and since the image of θ is precisely the set of all normal forms $t^{\pm 1}v_1t^{\pm 1}v_2 \dots$ with t -exponent sum divisible by k , the image of θ is precisely K .

Next, let Δ_* be Δ_k with an additional vertex $*$ and k additional edges $(i, *)$ ($i \in \mathbb{Z}/k$). Taking vertex groups to be the V_i and V_* (corresponding to $*$), edge groups to be the E_i and copies of the V_i (corresponding to $(i, *)$), and attaching homomorphisms to be the f_i and g_i , along with the identity on V_i (corresponding to the initial vertex of $(i, *)$) and h_i (corresponding to the terminal vertex of $(i, *)$), we get a graph of groups $(\{V_*, V_i, E_i\}, \Delta_*)$.

Since $(\{V_i, E_i\}, \Delta_k)$ is a sub-(graph of groups) of $(\{V_*, V_i, E_i\}, \Delta_*)$, by the normal form theorem, its fundamental group K is a subgroup of the fundamental group $\pi_1(\{V_*, V_i, E_i\}, \Delta_*)$. Furthermore, if T_Δ is the spanning tree of Δ_* whose edges are the $(i, *)$ and T_Γ is the (unique) spanning tree of Γ_k , by (4)–(5), the group $\pi_1(\{V_*, V_i, E_i\}, \Delta_*, T_\Delta)$ is isomorphic to $\pi_1(\{V_*, E_i\}, \Gamma_k, T_\Gamma)$. The lemma follows. $\blacksquare$

For the rest of this section, we let $S, m, \phi : S' \rightarrow S''$, and $G(S, m, \phi)$ be as defined in Section 1. Let Λ be the (directed) graph with vertex set S and an edge $(s, \phi(s))$ for each $s \in S'$. Since every vertex of Λ has in-degree and out-degree each at most 1, Λ is a disjoint union of loops and (non-closed) paths. Choose an integer k to be a multiple of the length of each loop in Λ and strictly greater than twice the length of each path in Λ .

Let $\overline{S}$ be the quotient of the direct product $\mathbb{Z}/k \times S$ by the equivalence relation generated by $(i, s) \sim (i+1, \phi(s))$ for all $i \in \mathbb{Z}/k$ and $s \in S'$. More geometrically, let $\widehat{\Lambda}$ be the (directed) graph with vertex set $\mathbb{Z}/k \times S$ and an edge $((i, s), (i+1, \phi(s)))$ for each $s \in S'$ and each $i \in \mathbb{Z}/k$. Then the map $(i, s) \mapsto s$ on vertex sets extends to a k -fold covering map $\pi : \widehat{\Lambda} \rightarrow \Lambda$, and each element of $\overline{S}$ is the vertex set of a connected component of $\widehat{\Lambda}$.

For $i \in \mathbb{Z}/k$, define $\psi_i : S \rightarrow \overline{S}$ by letting $\psi_i(s)$ be the equivalence class of (i, s) . We would like to define a labelling function $\overline{m}$ on $\overline{S}$ by

$$\overline{m}(x, y) = \begin{cases} m(s, t) & \text{if } x = \psi_i(s) \text{ and } y = \psi_i(t) \text{ for some } i \in \mathbb{Z}/k, \\ \infty & \text{otherwise.} \end{cases} \quad (6)$$

Lemma 5 *The labelling function $\overline{m}$ is well-defined.*

The proof of Lemma 5 uses a variation on an idea from [11, Lem. 5.7].

Proof. Let $p : (i, s) \mapsto i$ be the natural projection from $\widehat{\Lambda}$ to the graph Δ_k defined above, and let $\pi : \widehat{\Lambda} \rightarrow \Lambda$ be the covering map defined above. If $\hat{\ell}$ is a component of $\widehat{\Lambda}$ such that $\ell = \pi(\hat{\ell})$ is a closed loop, then since the length of ℓ divides k , p induces a graph isomorphism from $\hat{\ell}$ to Δ_k . Similarly, if $\hat{\ell}$ is a component of $\widehat{\Lambda}$ such that $\ell = \pi(\hat{\ell})$ is a non-closed path in Λ , then p induces a graph isomorphism from $\hat{\ell}$ to $p(\hat{\ell})$, a path of length strictly less than $k/2$. It

follows that for any two components $\hat{\ell}, \hat{\ell}'$ of $\hat{\Lambda}$, the intersection $p(\hat{\ell}) \cap p(\hat{\ell}')$ is a connected subgraph of Δ_k .

So now, suppose we have $i, j \in \mathbb{Z}/k$ and $s, s', t, t' \in S$ such that $x = \psi_i(s) = \psi_j(s')$ and $y = \psi_i(t) = \psi_j(t')$. Equivalently, suppose the vertices (i, s) and (j, s') lie in the same component x of $\hat{\Lambda}$ and the vertices (i, t) and (j, t') lie in the same component y of $\hat{\Lambda}$. Since p induces graph isomorphisms from x and y to their images $p(x)$ and $p(y)$, respectively, and $p(x) \cap p(y)$ is connected, for some $n \geq 0$, either both $(j, s') = (i + n, \phi^n(s))$ and $(j, t') = (i + n, \phi^n(t))$, or both $(i, s) = (j + n, \phi^n(s'))$ and $(i, t) = (j + n, \phi^n(t'))$. In either case, by (2), we see that $m(s, t) = m(s', t')$. ■

Proof of Theorem 1. Taking the Artin case first, let $V = A(S, m)$, $E = A(S', m)$, and $V_* = A(\overline{S}, \overline{m})$ (which is well-defined by Lemma 5). Also, let $f : E \rightarrow V$ be induced by the inclusion of S' in S and let $g : E \rightarrow V$ be induced by $\phi : S' \rightarrow S''$. Finally, since k is a multiple of the length of each loop in Λ and strictly longer than each path in Λ , every lift of a loop or path in Λ intersects each image $\psi_i(S)$ at most once, which means that each of the maps $\psi_i : S \rightarrow \overline{S}$ is injective and each ψ_i induces an injection $h_i : V \rightarrow V_*$. Applying Lemma 4 with this choice of V , E , V_* , f , g , and h_i , we see that $G(S, m, \phi)$ has a subgroup of index k that embeds in $G_* = \pi_1(\{V_*, E_i\}, \Gamma_k)$.

It remains to show that G_* is an Artin group. By (5), we see that G_* has a presentation with generators $\overline{S} \cup \{t_i\}$ ($i \in \mathbb{Z}/k$), subject to the Artin relations for $(\overline{S}, \overline{m})$ together with the relations

$$t_i^{-1} \psi_i(s) t_i = \psi_{i+1}(\phi(s)) \quad (7)$$

for all $i \in \mathbb{Z}/k$ and $s \in S'$. However, since $(i, s) = (i + 1, \phi(s))$ in $\overline{S}$, (7) becomes the Artin relation $\psi_i(s) t_i = t_i \psi_i(s)$. Therefore, if

$$m^+(x, y) = \begin{cases} \overline{m}(x, y) & \text{if } x, y \in \overline{S}, \\ 2 & \text{if } \{x, y\} = \{t_i, \psi_i(s)\} \text{ for some } i \in \mathbb{Z}/k, \\ \infty & \text{otherwise,} \end{cases} \quad (8)$$

we see that $G_* = A(\overline{S} \cup \{t_i\}, m^+)$.

Now consider the Coxeter case. Replacing $V = A(S, m)$, $E = A(S', m)$, and $V_* = A(\overline{S}, \overline{m})$ with $V = W(S, m)$, $E = W(S', m)$, and $V_* = W(\overline{S}, \overline{m})$, the previous proof carries through, except that G_* is not a Coxeter group. However, consider $W^+ = W(\overline{S} \cup \{u_i, u'_i\}, m^+)$, where

$$m^+(x, y) = \begin{cases} \overline{m}(x, y) & \text{if } x, y \in \overline{S}, \\ 2 & \text{if } \{x, y\} = \{u_i, \psi_i(s)\} \text{ for some } i \in \mathbb{Z}/k, \\ 2 & \text{if } \{x, y\} = \{u'_i, \psi_i(s)\} \text{ for some } i \in \mathbb{Z}/k, \\ \infty & \text{otherwise.} \end{cases} \quad (9)$$

It follows from Tits' solution to the word problem for Coxeter groups that the homomorphism $\theta : G_* \rightarrow W^+$ defined by $\theta(s) = s$ for $s \in \overline{S}$ and $\theta(t_i) = u_i u'_i$ is injective, as reduced (graph of groups) words in G_* are sent to reduced (Coxeter) words in W^+ . The theorem follows. ■

3 An open question

Can Theorem 1 be extended to “Artin graphs of Artin groups” (defined similarly to Artin HNN-extensions)? Note that since an “Artin amalgamated free product” is itself an Artin group, it is enough to consider multiple-Artin-HNN-extensions. In that case, the key point is to generalize Lemma 5, but it is not clear to us how (or if) that can be done.

One may of course also consider the analogous Coxeter question(s).

References

- [1] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapitres 4–6, Masson, Paris, 1981.
- [2] K. S. Brown, *Buildings*, Springer, New York, 1998.
- [3] M. W. Davis T. Januszkiewicz, ‘Right-angled Artin groups are commensurable with right-angled Coxeter groups’, *J. Pure Appl. Algebra* 153 (2000) 229–235.
- [4] A. Hatcher, *Algebraic Topology*, Cambridge Univ. Press, Cambridge, 2002.
- [5] T. Hsu D. T. Wise, ‘On linear and residual properties of graph products’, *Michigan Math. J.* 46 (1999), 251–259.
- [6] I. J. Leary, ‘On finite subgroups of groups of type VF’, *Geom. Topol.* 9 (2005) 1953–1976.
- [7] I. J. Leary B. E. A. Nucinkis, ‘Some groups of type VF’, *Invent. Math.* 151 (2003) 135–165.
- [8] H. Van der Lek, *The homotopy type of complex hyperplane complements*, Ph.D. thesis, Nijmegen, 1983.
- [9] L. Paris, ‘Parabolic subgroups of Artin groups’, *J. Algebra* 196 (1997) 369–399.
- [10] J.-P. Serre, *Trees*, Springer-Verlag, Berlin, 1980.
- [11] D. T. Wise, ‘Subgroup separability of the figure 8 knot group’, *Topology* 45 (2006) 421–463.

T. Hsu

Department of Mathematics San José State University San José CA 95192-0103
USA

hsu@math.sjsu.edu

I. J. Leary

Department of Mathematics The Ohio State University 231 W. 18th Avenue
Columbus OH 43210 USA

leary@math.ohio-state.edu